




Dams






Sub-standard Culverts







Culvert Problems

- Inlet or Outlet drop
- Physical barriers
- Debris accumulation
- Excessive velocities
- Insufficient water depth
- Flow contraction (turbulence)
- Orifice Flow
- Absence of bank edge areas
- Discontinuity of channel substrate







Ecological Processes


- Energy flow / Food chain support
- Interdependencies
- Predator-prey dynamics
- Hydrology
- Natural disturbances
- Movement of organisms and materials

Interdependencies



The occurrence of some species is dependent on the present of others. For example many freshwater mussel species are dependent on specific fish hosts to complete their life cycles.





Predator-Prey Dynamics

What are the impacts of

- fish stocking
- introduced fish
- absence of native anadromous fish
- ...on stream communities.



Importance of Movement

- Daily movements
- Changes in habitat conditions
- Reproduction
- Exploit vacant habitat
- Population continuity
- Dispersal

Adult Spawning Migrations





Spawning Habitat Eggs & Alevin

Requirements:

• Clean, well oxygenated gravels



Salmon Fry

Requirements:

- Margin habitats with slow-moderate current
- Sufficient invertebrate prey
- Interstitial spaces





Early Parr

Requirements:

- Habitat with moderate-swift current
- Sufficient invertebrate prey
- Adequate interstitial spaces






Late Parr / Pre-smolt

Requirements:

- Overwinter cover
- Require larger shelter
- Appropriate water chemistry
- Ability to emigrate from natal streams at certain times of the year



At this stage salmon make extensive movements seeking appropriate winter habitat



Habitat suitable for Fallfish

Site4.shp



(



1


Impacts of River & Stream Crossings


- Habitat loss and degradation
- Roadkill leading to loss of populations
- Alteration of Ecological Processes
- Reduced access to vital habitats
- Population fragmentation & isolation
- Disruption of processes that maintain regional populations





Reduced Access to Vital Habitats


- Spawning habitat
- Nursery habitat
- Foraging areas
- Deep water refuges
- Seasonal habitats











Population Fragmentation and Isolation

- Barriers to movement subdivide or isolate populations
- Smaller and more isolated populations are more vulnerable to:
 - extinction due to chance events
 - genetic changes

Population Viability

Short-term viability

$$N_e = 50 \text{ to } 200 +$$

Long-term viability

$$N_e = 500 \text{ to } 5000 +$$

Processes that Maintain Regional Populations ("Metapopulations")

- Supplementation ("rescue effect")
- Gene flow
- Re-colonization



Mudpuppy

Amphiuma



Hellbender



Cope's Giant Salamander



Softshell turtles

Musk turtles










Importance of Small Streams


- Make up a large percentage of stream miles
- Cumulatively provide more habitat than large rivers
- Support species not found in larger streams and rivers
- Provide important spawning & nursery habitat for fish























Aldo Leopold, 1953:

"If the biota, in the course of eons, has built something we like but do not understand, then who but a fool would discard seemingly useless parts? To keep every cog and wheel is the first precaution of intelligent tinkering."